

Level of IL-6 predicts respiratory failure in hospitalized symptomatic COVID-19 patients

Tobias Herold MD^{1,2,3,4*}, Vindi Jurinovic PhD^{1,5*}, Chiara Annreich^{1,4}, Johannes C. Hellmuth MD¹, Michael von Bergwelt-Baildon MD PhD^{1,3,6}, Matthias Klein MD^{2,7} and Tobias Weinberger MD^{2,8}

1 Department of Medicine III, University Hospital, LMU Munich, Munich, Germany

2 Emergency Department, University Hospital, LMU Munich, Munich, Germany

3 Nine-i Investigator

4 Research Unit Apoptosis in Hematopoietic Stem Cells, Helmholtz Zentrum München, German Center for Environmental Health (HMGU), Munich, Germany

5 Institute for Medical Information Processing, Biometry and Epidemiology, LMU Munich, Munich, Germany

6 Intensive Care in Hematologic and Oncologic Patients (iCHOP), Cologne, Germany

7 Department of Neurology, University Hospital, LMU Munich, Munich, Germany

8 Department of Medicine I, University Hospital, LMU Munich, Munich, Germany

* Contributed equally

Corresponding Author:

Tobias Herold, MD; Marchioninstr. 15; 81377 Munich; Germany; Phone: +49 89 4400-0; FAX: +49 89 4400-74242; Email: tobias.herold@med.uni-muenchen.de

Conflict of interest: The authors declare no conflict of interest.

Abstract

The pandemic Coronavirus-disease 19 (COVID-19) is characterized by a heterogeneous clinical course. While most patients experience only mild symptoms, a relevant proportion develop severe disease progression with increasing hypoxia up to acute respiratory distress syndrome. The substantial number of patients with severe disease have strained intensive care capacities to an unprecedented level. Owing to the highly variable course and lack of reliable predictors for deterioration, we aimed to identify variables that allow the prediction of patients with a high risk of respiratory failure and need of mechanical ventilation

Patients with PCR proven symptomatic COVID-19 infection hospitalized at our institution from 29th February to 27th March 2020 (n=40) were analyzed for baseline clinical and laboratory findings. Patients requiring mechanical ventilation 13/40 (32.5%) did not differ in age, comorbidities, radiological findings, respiratory rate or qSofa score. However, elevated interleukin-6 (IL-6) was strongly associated with the need for mechanical ventilation ($p=1.2 \cdot 10^{-5}$). In addition, the maximal IL-6 level (cutoff 80 pg/ml) for each patient during disease predicted respiratory failure with high accuracy ($p=1.7 \cdot 10^{-8}$, AUC=0.98). The risk of respiratory failure for patients with IL-6 levels of ≥ 80 pg /ml was 22 times higher compared to patients with lower IL-6 levels.

In the current situation with overwhelmed intensive care units and overcrowded emergency rooms, correct triage of patients in need of intensive care is crucial. Our study shows that IL-6 is an effective marker that might be able to predict upcoming respiratory failure with high accuracy and help physicians correctly allocate patients at an early stage.

Introduction

The pandemic Coronavirus-disease 19 (COVID-19) is characterized clinically by a highly variable course. While most patients experience only mild symptoms, a relevant proportion develop severe disease progression with increasing hypoxia up to acute respiratory distress syndrome. About 5% of patients require intensive care treatment including mechanical ventilation.¹⁻³ This variability of COVID-19 and the shortage of health care resources in heavily affected regions make efficient allocation of resources towards patients at high risk for deterioration crucial.⁴ We aimed to identify variables that allow the prediction of patients with a high risk of respiratory failure and need of mechanical ventilation.

Methods

All patients with PCR proven symptomatic COVID-19 infection hospitalized at our institution from 29th February to 27th March 2020 (n=40) were analyzed for baseline clinical and laboratory findings. Decision on endotracheal intubation was made following internationally accepted recommendations (PaO₂/FiO₂ <150mmHg).⁵ Patient data were anonymized for analysis and the study was approved by the local ethics committee (No: 20-245). All variables with less than 50% of missing data were tested for the association with respiratory failure. Categorical variables were compared with the χ^2 -test, and numerical variables were tested with the Mann-Whitney-U-Test. The p-values were adjusted for multiple testing with the Bonferroni-Holm-Method. An adjusted p-value (q-value) of ≤ 0.05 was considered significant.

Results

In total, 13/40 (32.5%) patients deteriorated during hospitalization and required mechanical ventilation. The time from hospital admission to intubation varied from less than one hour to 9 days (median 2 days). All patients who required intubation were of male sex, compared to 59% of males in the group that did not require intubation (p=0.020, q=0.057). Patients requiring mechanical ventilation did not differ in age, comorbidities, radiological findings, respiratory rate or qSofa score (table).

Pulse, markers of inflammation, LDH and creatinine at admission were associated with respiratory failure (table).

Elevated interleukin-6 (IL-6) was very strongly associated with the need for mechanical ventilation (figure 1A, $p=1.2 \cdot 10^{-5}$, $q=0.00032$).

In addition to IL-6 values at first assessment, follow-up data on IL-6 were also available (median number of values per patient: 6, range 1-10). These data were used to assess the maximal IL-6 level for each patient during disease (for patients requiring ventilation, only values before intubation were used). These values predicted respiratory failure with high accuracy (figure 1B/C, $p=1.7 \cdot 10^{-8}$, $AUC=0.98$). The statistically optimal cutoff for IL-6 was 80 pg/ml, identifying all but one patient with respiratory failure correctly, and misclassifying only one patient that still does not require intubation. The risk of respiratory failure for patients with IL-6 levels of ≥ 80 pg/ml was 92% and thus 22 times higher compared to patients with lower IL-6 levels. After reaching an IL-6 value of 80pg/ml, the median time to mechanical ventilation was 1.5 days (range 0–4 days).

Discussion

Even though IL-6 levels were significantly elevated in patients requiring ventilation, they are relatively low compared to levels observed in patients with septic shock.⁶ Our data suggests that even moderately elevated IL-6 levels above 80pg/ml are sufficient to identify COVID-19 patients with a high risk of respiratory failure. Further studies and larger sample sizes will be needed to validate our findings and possibly determine a more accurate cutoff. To date, it is unclear whether IL-6 merely represents a biomarker or a central pathogenetic element of severe COVID-19 that should be used as a parameter for therapeutic intervention.

In the current situation with overwhelmed intensive care units and overcrowded emergency rooms, correct triage of patients in need of intensive care is crucial. Our study shows that IL-6 is an effective marker that might be able to predict upcoming respiratory failure with high accuracy and help physicians correctly allocate patients at an early stage.

Acknowledgement: We thank Oliver T. Keppler, the Task Force Corona at the University Hospital, LMU Munich and all CORKUM investigators as well as all health care workers for their outstanding service.

References

1. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020. doi:10.1016/S0140-6736(20)30566-3.
2. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. 2020. doi:10.1001/jama.2020.1585.
3. Guan W-J, Ni Z-Y, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020. doi:10.1056/NEJMoa2002032.
4. Emanuel EJ, Persad G, Upshur R, et al. Fair Allocation of Scarce Medical Resources in the Time of Covid-19. *N Engl J Med*. 2020. doi:10.1056/NEJMs2005114.
5. Bellani G, Laffey JG, Pham T, et al. Epidemiology, Patterns of Care, and Mortality for Patients With Acute Respiratory Distress Syndrome in Intensive Care Units in 50 Countries. *JAMA*. 2016;315(8):788-800. doi:10.1001/jama.2016.0291.
6. Surbatovic M, Popovic N, Vojvodic D, et al. Cytokine profile in severe Gram-positive and Gram-negative abdominal sepsis. *Sci Rep*. 2015;5:11355. doi:10.1038/srep11355.

Table

Variable	Evaluable	Median (range) / n (%)	Mechanical ventilation		p-value	q-value
			No (n = 27)	Yes (n = 13)		
Baseline Characteristics						
Age (years)	40	57 (19 - 81)	54 (19 - 80)	64 (45 - 81)	0.15	0.28
Respiratory rate (/min)	33	18 (14 - 40)	18 (14 - 32)	23 (15 - 40)	0.075	0.15
Pulse (/min)	33	82 (54 - 130)	77 (54 - 111)	95 (80 - 130)	0.0028	0.012
Male gender	40	29 (72)	16 (59)	13 (100)	0.020	0.058
Any comorbidities	39	32 (82)	20 (77)	12 (92)	0.46	0.70
Hypertension	36	19 (53)	10 (40)	9 (82)	0.051	0.11
Diabetes mellitus	37	3 (8)	1 (4)	2 (17)	0.50	0.72
Coronary heart disease	34	3 (9)	3 (12)	0 (0)	0.61	0.80
Chronic obstructive lung disease	35	3 (9)	2 (8)	1 (9)	>0.99	>0.99
Computed Tomography						
Consolidation	35	20 (57)	14 (61)	6 (50)	0.80	0.99
Ground glass opacity	35	31 (89)	20 (87)	11 (92)	>0.99	>0.99
Bilateral infiltration	35	31 (89)	21 (91)	10 (83)	0.89	>0.99
Scores						
qSOFA score	31	12 (39)	7 (32)	5 (56)	0.41	0.66
CURB-65 score ≥ 1	23	7 (30)	5 (31)	2 (29)	>0.99	>0.99
Laboratory parameters	Evaluable	Median (range)	Mechanical ventilation		p-value	q-value
			No (n = 27)	Yes (n = 13)		
Lymphocyte count (%)	36	19 (4 - 45)	21 (4 - 45)	15 (6 - 26)	0.050	0.11
CRP (mg/dl)	40	2.8 (0 - 31.5)	1.7 (0 - 31.5)	7.8 (1.6 - 17.1)	0.0019	0.012
Bilirubin (mg/dl)	36	0.5 (0.2 - 1.9)	0.5 (0.2 - 1.2)	0.5 (0.4 - 1.9)	0.93	>0.99
WBC (G/l)	40	5.295 (2.12 - 308)	4.75 (2.12 - 12.5)	6.64 (4.99 - 308)	0.0014	0.012
LDH (U/l)	38	292 (182 - 1078)	281 (182 - 619)	346 (252 - 1078)	0.0026	0.012
PCT (ng/ml)	37	0 (0 - 5)	0 (0 - 0.6)	0.1 (0 - 5)	0.011	0.041
IL6 (pg/ml)	37	27.1 (0 - 430)	19.6 (0 - 76.5)	121 (19.2 - 430)	0.000012	0.00032
Thrombocyte count (G/l)	40	165 (88 - 440)	186 (88 - 334)	160 (1 - 440)	0.59	0.80
Troponin T (ng/ml)	34	0 (0 - 0.032)	0 (0 - 0.022)	0 (0 - 0.032)	0.018	0.058
Creatinine (mg/dl)	40	0.9 (0.4 - 2.1)	0.9 (0.4 - 1.3)	1.0 (0.9 - 2.1)	0.00034	0.0044
D-Dimer	30	0.7 (0 - 2.9)	0.6 (0 - 2.2)	1.1 (0.6 - 2.9)	0.028	0.074
Ferritin (ng/ml)	27	644 (64 - 2153)	606 (64 - 1748)	810 (431 - 2153)	0.16	0.28

CRP = C-Reactive Protein; WBC= White blood cell count; LDH = Lactate Dehydrogenase;
PCT = Procalcitonin; IL6 = Interleukin-6

Figure legend

Figure 1: Box plot showing IL-6 levels at first assessment $p=1.2 \cdot 10^{-5}$ (A) and maximal IL-6 levels before mechanical ventilation $p=1.7 \cdot 10^{-8}$ (B). Receiver operating characteristic (ROC) curve of maximal IL-6 levels before mechanical ventilation, $AUC=0.98$ (C).

Figure 1

